Author Index

Abatangelo, G., see Brun, P. 530
Abe, J.
—, Asada, A., Fujimori, M., Imaoka, S. and Funae, Y.
Binding of lidocaine to plasma proteins resolved by high-performance liquid chromatography 562
Ageta, T., see Mizobuchi, N. 550
Akiyama, H., see Koike, M. 235
Alebić-Kolbah, T.
— and Hiršl-Starčević, S.
Determination of piracetam in serum by gas chromatography 556
Amiss, T.J.
—, Tyczkowska, K.L. and Aucoin, D.P.
Analysis of taurine in feline plasma and whole blood by liquid chromatography with fluorimetric detection and confirmation by thermospray mass spectrometry 375
Annan, R.S.
—, Kim, C. and Martyn, J.
Measurement of D-tubocurarine chloride in human urine using solid-phase extraction and reversed-phase high-performance liquid chromatography with ultraviolet detection 228
Anttila, M., see Laakso, S. 475
Argoudelis, C.J.
Simple isocratic high-performance liquid chromatographic method for measuring pyridoxine kinase activity in crude biological extracts 25
Asada, A., see Abe, J., 562
Asai, S., see Imamura, Y. 282
Aucin, D.P., see Amiss, T.J. 375
Awni, W.M., see Maloney, J.A. 487
Baars, A.M., see Vree, T.B. 581
Badet, J., see Jacquot Dourges, M.A. 35
Balis, F.M., see Carpen, M.E. 69
Barritault, D., see Jacquot Dourges, M.A. 35
Bayne, W.F., see Matuszewski, B.K. 461
Benakis, A., see Schopfer, C. 223
Beneken Kolmer, E.W.J., see Vree, T.B. 119
Blanchard, J.R.T.
— and Tasker, R.A.R.
High-performance liquid chromatographic assay for domoic acid in serum of different species 546
Boelsterli, U.A., see Bouis, P. 447
Bosch, R., see Vree, T.B. 119
Bouis, P.
—, Taccard, G. and Boelsterli, U.A.
Determination of cocaine and norcocaine in plasma and cell cultures using high performance liquid chromatography 447
Bradwin, G.R., see Underwood, R.H. 180
Brandin, S., see Polacek, J. 151
Brun, P.
—, De Galateo, A., Camporese, A., Cortivo, R. and Abatangelo, G.
Analysis of hyaluronic acid in synovial fluid by reversed-phase liquid chromatography 530
Bruns, M.B., see Lakings, D.B. 273
Bullen, W.W.
—, Chang, T. and Whitfield, L.R.
High-performance liquid chromatographic assay for trimetrexate in human plasma 266
Burghardt, C., see Janero, D.R. 11
Burton, L.C.
—, Loftus, N.J., Vere, D.W. and Whelpton, R.
Determination of plasma nefopam by liquid chromatography and electrochemical detection 159
Camporese, A., see Brun, P. 530
Candito, M.
—, Krstulovic, A.M., Sbirazzuoli, V. and Chambon, P.
Proposal for the standardization of the calibration method for the assay of plasma catecholamines 194
Cannell, H., see Whelpton, R. 215
Carpen, M.E.
—, Poplack, D.G., Pizzo, P.A. and Balis, F.M.
High-performance liquid chromatographic method for analysis of 2′,3′-dideoxyinosine in human body fluids 69
Chambon, P., see Candito, M. 194
Chang, D., see Xie, F. 109
Chang, T., see Bullen, W.W. 266
Chao, W.-F.
—, Kai, M. and Ohkura, Y.
High-performance liquid chromato-

graphic determination of a cysteine protease inhibitor and its ethyl ester in mouse serum and muscle by pre-column fluorescence derivatization 77
Charleson, S., see Nicoll-Griffith, D. 341
Chow, D., see Lau, S.W.J. 87
Coetzee, J., see Uebel, R.A. 293
Conner, M.K., see Mayernik, J.A. 407
Constanzer, M.L., see Matuszewski, B.K. 461
Cortivo, R., see Brun, P. 530
Courty, J., see Jacquot Dourges, M.A. 35
Culling-Berlund, A.J., see Newcomb, S.A. 47
Damm, P., see Lehr, K.H. 497
Dasgupta, A.
—, Rosenzweig, I.B., Turgeon, J. and Raisys, V.A.
Encainide and metabolites analysis in serum or plasma using a reversed-phase high-performance liquid chromatographic technique 260
Davis, T.P., see Newcomb, S.A. 47
De Galateo, A., see Brun, P. 530
De Mey, C., see Meineke, I. 240
De Schepper, P.J., see Verbesselt, R. 515
Ding, W.
—, Veening, H. and Van Effen, R.M.
Determination of guanidino compounds by anion-exchange chromatography and amperometric detection 355
Dourges, M.A., Jacquot, see Jacquot Dourges, M.A. 35
Dudson, P., see Whelpton, R. 215
Echizen, H.
—, Nakura, M. and Ishizaki, I.
Rapid and simple high-performance liquid chromatographic determination of saliva antipyrine for routine antipyrine test 296
Effen, R.M. van, see Ding, W. 355
Egloff, M., see Frémont, S. 186
Eguchi, K., see Morita, I. 367
Embree, L.
— and McErlane, K.M.
Electrochemical detection of the 3,5-dinitrobenzoyl derivative of digoxin by high-performance liquid chromatography 439
Ethier, D., see Nicoll-Griffith, D. 341
Felden, F., see Frémont, S. 186
Feldman, S., see Lau, S.WJ. 87
Foster, R.T., see Piquette-Miller, M. 129
Franklin, M.
Determination of plasma buspirone by high-performance liquid chromatography with coulometric detection 590
Frémont, S.
—, Guéant, J.-L., Felden, F., Egloff, M., Gérard, A. and Nicolas, J.-P.
Dihydrotestosterone binding capacity of androgen-binding protein in tissue extract using high-performance size exclusion chromatography 186
Friss, J.M., see Lakings, D.B. 273
Fujimori, M., see Abe, J. 562
Funae, Y., see Abe, J. 562
Galateo, A. De, see Brun, P. 530
Gérard, A., see Frémont, S. 186
Giam, C.S., see Mayernik, J.A. 407
Gotoh, Y., see Imamura, Y. 282
Gould, D.H., see Savage, J.C. 540
Guéant, J.-L., see Frémont, S. 186
Gulino, D., see Jacquot Dourges, M.A. 35
Gustafsson, B., see Polacek, J. 151
Hakusui, H., see Nakaoka, M. 254
Harimoto, T., see Nishioka, R. 210
Hartnack, S., see Meineke, I. 240
Hatta, Y., see Tomita, M. 174
Hattori, M., see Morita, I. 367
Hawtrey, A.O., see Uebel, R.A. 293
Hekster, Y.A., see Vree, T.B. 581
Hessey, G.A., see Matuszewski, B.K. 461
Hill, H.M., see Selinger, K. 597
Hirokami, M., see Matsumoto, M. 1
Hiršl-Starčević, S., see Alebić-Kolbah, T. 556
Hocke, G.
—, Kaffarnik, H., Münscher, G. and Steinmetz, A.
Purification of human serum amyloid A by anion-exchange fast protein liquid chromatography 203
Hooper, W.D., see Treston, A.M. 59
Hori, T., see Tatsuhara, T. 507
Hsieh, Y., see Xie, F. 109
Imamura, Y.
—, Shibanoki, S., Kubo, T., Kogure, M., Ishii, Y., Gotoh, Y., Asai, S. and Ishikawa, K.
High-performance liquid chromatographic determination of *cis*-N-(1-benzyl-2-methylpyrrolidin-3-yl)-5-chloro-2-methoxy-4-methylaminobenzamide in rat blood and brain using electrochemical detection 282

Imaoka, S., see Abe, J. 562
Imazawa, M., see Shimamura, M. 535
Ishii, Y., see Imamura, Y. 282
Ishikawa, K., see Imamura, Y. 282
Ishizaki, I., see Echizen, H. 296
Itakura, K., see Verbesselt, R. 515
Ito, H.
—, Yamamoto, H., Kimura, Y., Kambe, H., Okochi, T. and Kishimoto, S.
Affinity chromatography of human plasma gelsolin with polyphosphate compounds on immobilized Cibacron Blue F3GA 397
Jacquot Dourges, M.A.
—, Gulino, D., Courty, J., Badet, J., Barritault, D. and Jozefonvicz, J.
Affinity chromatography of fibroblast growth factors on substituted polystyrene 35
Jamali, F., see Piquette-Miller, M. 129
Janero, D.R.
— and Burghardt, C.
Solid-phase extraction on silica cartidges as an aid to platelet-activating factor enrichment and analysis 11
Janssen, M.G.A., see Wuis, E.W. 575
Jiang, J., see Xie, F. 109
Jozefonvicz, J., see Jacquot Dourges, M.A. 35
Kaffarnik, H., see Hocke, G. 203
Kai, M., see Chao, W.-F. 77
Kambe, H., see Ito, H. 397
Kanemitsu, F.
— and Okigaki, T.
Clinical application of subforms of creatine kinase MM and macro creatine kinases 423
Kargas, G.
—, Rudy, T., Spennetta, T., Takayama, K., Querishi, N. and Shrago, E.
Separation and quantitation of long-chain free fatty acids in human serum by high-performance liquid chromatography 331
Kawai, S., see Tomita, M. 174
Kawamoto, M., see Morita, I. 367
Kim, C., see Annan, R.S. 228
Kimura, Y., see Ito, H. 397
Kishimoto, S., see Ito, H. 397
Kleijn, E. van der, see Wuis, E.W. 575
Knapp, S.
—, Wardlow, M.L. and Thal, L.J.
Sensitive analysis of plasma physostigmine levels using dual-cell electrochemistry in the redox mode 97
Knörr, W.
— and Spiteller, G.
Simple method for the analysis of glycerol enol ethers derived from plasmalogens in complex lipid mixtures and subsequent determination of the aldehydic components by gas chromatography–mass spectrometry 303
Kodama, H., see Mizobuchi, N. 550
Kogure, M., see Imamura, Y. 282
Koike, M.
—, Akiyama, H. and Shimizu, T.
High-performance liquid chromatographic procedure for the determination of rat plasma concentrations of a new antibacterial agent, (±)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidyl)-5-methyl-1-oxo-1*H*,5*H*-benzo[*i*,*j*]quinolizine-2-carboxylic acid, for topical use 235
Kolmer, E.W.J. Beneken, see Vree, T.B. 119
Krstulovic, A.M., see Candito, M. 194
Kubo, T., see Imamura, Y. 282
Laakso, S.
—, Scheinin, M. and Anttila, M.
Determination of erythromycin base and 2′-acetylerythromycin in human plasma using high-performance liquid chromatography with electrochemical detection 475
Lakings, D.B.
—, Friis, J.M. and Bruns, M.B.
Determination of ditekiren, a renin inhibitor peptide, in monkey serum using high-performance liquid chromatography with solid-phase extraction 273
Lau, S.W.J.
—, Low, D. and Feldman, S.
Simultaneous determination of physostigmine and tetrahydroaminoacridine in a transdermal permeation study by high-performance liquid chromatography 87
Lehr, K.H.
— and Damm, P.
Simultaneous determination of the sulphonylurea glimepiride and its metabolites in human serum and urine by high-performance liquid chromatography after pre-column derivatization 497
Leis, H.-J.
—, Welz, W. and Malle, E.
11-Dehydrothromboxane B_2: convenient preparation of its stable isotope-labelled

analogue for use in quantitative gas chromatography–mass spectrometry 169
—, see Welz, W. 319
Lhermitte, M., see Pommery, J. 569
Li, J., see Xie, F. 109
Loftus, N.J., see Burton, L.C. 159
McErlane, K.M., see Embree, L. 439
Malle, E., see Leis, H.-J. 169
—, see Welz, W. 319
Maloney, J.A.
— and Awni, W.M.
High-performance liquid chromatographic determination of isepamicin in plasma, urine and dialysate 487
Martea, M., see Vree, T.B. 119
Martin, S.G., see Stratford, M.R.L. 383
Martyn, J., see Annan, R.S. 228
Mathieu, D., see Pommery, J. 569
Mathieu, M., see Pommery, J. 569
Matsumoto, M.
—, Togashi, H., Yoshioka, M., Hirokami, M., Morii, K. and Saito, H.
Simultaneous high-performance liquid chromatographic determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaesthetized normotensive rats 1
Matuszewski, B.K.
—, Constanzer, M.L., Hessey, G.A. and Bayne, W.F.
Development of direct stereoselective and non-stereoselective assays in biological fluids for the enantiomers of a thieno[2,3-*b*]thiopyran-2-sulphonamide, a topically effective carbonic anhydrase inhibitor 461
Mayernik, J.A.
—, Conner, M.K. and Giam, C.S.
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Meineke, I.
—, Hartnack, S. and De Mey, C.
Determination of mepindolol in plasma after transdermal drug delivery by high-performance liquid chromatography with electrochemical detection 240
Mey, C. De, see Meineke, I. 240
Miyamoto, K., see Shimamura, M. 535
Mizobuchi, N.
—, Ageta, T. and Kodama, H.
Determination of N-monoacetylcystathionine in biological samples using isotachophoresis 550
Moore, T.J., see Underwood, R.H. 180
Morii, K., see Matsumoto, M. 1
Morita, I.
—, Kawamoto, M., Hattori, M., Eguchi, K., Sekiba, K. and Yoshida, H.
Determination of tryptophan and its metabolites in human plasma and serum by high-performance liquid chromatography with automated sample clean-up system 367
Münscher, G., see Hocke, G. 203
Murray, J.C., see Stratford, M.R.L. 383
Nakaoka, M.
— and Hakusui, H.
Measurement of the formation of a thioglucuronide, a metabolite of malotilate, in rat hepatic microsomes by high-performance liquid chromatography 254
Nakura, M., see Echizen, H. 296
Newcomb, S.A.
—, Culling-Berlund, A.J. and Davis, T.P.
Endogenous levels of β-carotene in human buccal mucosa cells by reversed-phase high-performance liquid chromatography 47
Nicolas, J.-P., see Frémont, S. 186
Nicoll-Griffith, D.
—, Zamboni, R., Rasmussen, J.B., Ethier, D., Charleson, S. and Tagari, P.
BIO-Fully Automated Sample Treatment high-performance liquid chromatography and radioimmunoassay for leukotriene E_4 in human urine from asthmatics 341
Nilsson, L.B.
Determination of remoxipride in plasma and urine by reversed-phase column liquid chromatography 139
Nishioka, R.
—, Harimoto, T., Umeda, I., Yamamoto, S. and Ōi, N.
Improved procedure for determination of indomethacin in plasma by capillary gas chromatography after solid-phase extraction 210
Ōhara, T., see Shimamura, M. 535
Ohkura, Y., see Chao, W.-F. 77
Ōi, N., see Nishioka, R. 210
Okigaki, T., see Kanemitsu, F. 423
Okochi, T., see Ito, H. 397

Okuyama, T., see Tomita, M. 174
Oliw, E.H.
Resolution of enantiomers of 19-hydroxyeicosatetraenoate and 18-hydroxyeicosatetraenoate by chiral phase high-performance liquid chromatography of naphthoyl ester derivatives 525
Ottersgård-Brorsson, A., see Polacek, J. 151
Pasutto, F.M., see Piquette-Miller, M. 129
Piquette-Miller, M.
—, Foster, R.T., Pasutto, F.M. and Jamali, F.
Stereospecific high-performance liquid chromatographic assay of acebutolol in human plasma and urine 129
Pizzo, P.A., see Carpen, M.E. 69
Polacek, J.
—, Gustafsson, B., Brandin, S. and Ottersgård-Brorsson, A.
Determination of tauromustine and its demethylated metabolites in plasma and urine 151
Pommery, J.
—, Mathieu, M., Mathieu, D. and Lhermitte, M.
High-performance liquid chromatographic determination of atrazine in human plasma 569
Poplack, D.G., see Carpen, M.E. 69
Prevost, J., see Selinger, K. 597
Querishi, N., see Kargas, G. 331
Raisys, V.A., see Dasgupta, A. 260
Rasmussen, J.B., see Nicoll-Griffith, D. 341
Rosenzweig, I.B., see Dasgupta, A. 260
Rudy, T., see Kargas, G. 331
Rustum, A.M.
Measurement of ibuprofen in human whole blood by reversed-phase ion-paired high-performance liquid chromatography using a pH-stable polymeric column 246
Saito, H., see Matsumoto, M. 1
Sattler, W., see Welz, W. 319
Savage, J.C.
— and Gould, D.H.
Determination of sulfide in brain tissue and rumen fluid by ion-interaction reversed-phase high-performance liquid chromatography 540
Sbirazzuoli, V., see Candito, M. 194
Scheinin, M., see Laakso, S. 475
Schepper, P.J. De, see Verbesselt, R. 515
Schopfer, C.
— and Benakis, A.
Simplified method for the determination of atracurium and laudanosine in pig plasma by high-performance liquid chromatography and fluorimetric detection 223
Sekiba, K., see Morita, I. 367
Selinger, K.
—, Prevost, J. and Hill, H.M.
High-performance liquid chromatography method for the determination of diphenhydramine in human plasma 597
Shibanoki, S., see Imamura, Y. 282
Shimamura, M.
—, Ōhara, T., Imazawa, M. and Miyamoto, K.
Purification of myelin-associated glycoprotein from calf brain using high performance liquid chromatography 535
Shimizu, M., see Tanaka, M. 289
Shimizu, T., see Koike, M. 235
Shimoda, M., see Vree, T.B. 119
Shrago, E., see Kargas, G. 331
Shu, H., see Xie, F. 109
Spennetta, T., see Kargas, G. 331
Spiteller, G., see Knörr, W. 303
Steegers-Theunissen, R.P.M., see Vree, T.B. 581
Steinmetz, A., see Hocke, G. 203
Stratford, M.R.L.
—, Watfa, R.R., Murray, J.C. and Martin, S.G.
Determination of collagen and protein turnover by high-performance liquid chromatography 383
Sugiro, T., see Verbesselt, R. 515
Tabuchi, F., see Tatsuhara, T. 507
Taccard, G., see Bouis, P. 447
Tagari, P., see Nicoll-Griffith, D. 341
Takayama, K., see Kargas, G. 331
Tanaka, M.
—, Shimizu, M. and Tsujimoto, Y.
High-performance liquid chromatographic determination of urinary mercapturic acids in rat 289
Tasker, R.A.R., see Blanchard, J.R.T. 546
Tatsuhara, T.
—, Tabuchi, F., Yamane, M. and Hori, T.
Rapid and simple method for the determination of nimustine hydrochloride in human blood and brain by high-performance liquid chromatography 507
Thal, L.J., see Knapp, S. 97
Tjandra-Maga, T.B., see Verbesselt, R. 515

Togashi, M., see Matsumoto, M. 1
Tomita, M.
—, Okuyama, T., Hatta, Y. and Kawai, S.
Determination of free malonaldehyde by gas chromatography with an electron-capture detector 174
Treston, A.M.
— and Hooper, W.D.
Metabolic studies with phenobarbitone, primidone and their N-alkyl derivatives: quantification of substrates and metabolites using chemical ionization gas chromatography–mass spectrometry 59
Tsujimoto, Y., see Tanaka, M. 289
Turgeon, J., see Dasgupta, A. 260
Tyczkowska, K.L., see Amiss, T.J. 375
Uebel, R.A.
—, Wium, C.A., Hawtrey, A.O. and Coetzee, J.
Electrochemical determination of 2,6-diisopropylphenol after high-performance liquid chromatography of extracts from serum 293
Umeda, I., see Nishioka, R. 210
Underwood, R.H.
—, Bradwin, G.R., Moore, T.J. and Williams, G.H.
Semi-automated high-performance liquid chromatographic method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test 180
Van der Kleijn, E., see Wuis, E.W. 575
Van Effen, R.M., see Ding, W. 355
Veening, H., see Ding, W. 355
Verbesselt, R.
—, Tjandra-Maga, T.B., De Schepper, P.J., Itakura, K. and Sugiro, T.
Determination of a platelet activating factor antagonist (CV-3988): methodology and clinical application 515
Vere, D.W., see Burton, L.C. 159
Vree, T.B., see Wuis, E.W. 575
—, Beneken Kolmer, E.W.J., Martea, M., Bosch, R. and Shimoda, M.
High-performance liquid chromatography of sulphadimethoxine and its N_1-glucuronide, N_4-acetyl and N_4-acetyl-N_1-glucuronide metabolites in human plasma and urine 119
—, Steegers-Theunissen, R.P.M., Baars, A.M. and Hekster, Y.A.
Direct high-performance liquid chromatographic analysis of *p*-hydroxyphenylphenylhydantoin glucuronide, the final metabolite of phenytoin, in human serum and urine 581
Wang, H., see Xie, F. 109
Wardlow, M.L., see Knapp, S. 97
Watfa, R.R., see Stratford, M.R.L. 383
Webster, K., see Whelpton, R. 215
Welz, W., see Leis, H.-J. 169
—, Sattler, W., Leis, H.-J. and Malle, E.
Rapid analysis of non-esterified fatty acids as methyl esters from different biological specimens by gas chromatography after one-step esterification 319
Whelpton, R., see Burton, L.C. 159
—, Dudson, P., Cannell, H. and Webster, K.
Determination of prilocaine in human plasma samples using high-performance liquid chromatography with dual-electrode electrochemical detection 215
Whitfield, L.R., see Bullen, W.W. 266
Williams, G.H., see Underwood, R.H. 180
Wium, C.A., see Uebel, R.A. 293
Wuis, E.W.
—, Janssen, M.G.A., Vree, T.B. and Van der Kleijn, E.
Determination of a dantrolene metabolite, 5-(*p*-nitrophenyl)-2-furoic acid, in plasma and urine by high-performance liquid chromatography 575
Xie, F.
—, Wang, H., Shu, H., Li, J., Jiang, J., Chang, J. and Hsieh, Y.
Separation and characterization of the metabolic products of lappaconitine in rat urine by high-performance liquid chromatography 109
Yamamoto, H., see Ito, H. 397
Yamamoto, S., see Nishioka, R. 210
Yamane, M., see Tatsuhara, T. 507
Yoshida, H., see Morita, I. 367
Yoshioka, M., see Matsumoto, M. 1
Zamboni, R., see Nicoll-Griffith, D. 341

Subject Index

Acebutolol
Stereospecific HPLC assay of acebutolol in human plasma and urine 129
Acetylcholine
Simultaneous HPLC determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaesthetized normotensive rats 1
2′-Acetylerythromycin
Determination of erythromycin base and 2′-acetylerythromycin in human plasma using HPLC with electrochemical detection 475
N_4-Acetylsulphadimethoxine
HPLC of sulphadimethoxine and its N_1-glucuronide, N_4-acetyl and N_4-acetyl-N_1-glucuronide metabolites in human plasma and urine 119
N_4-Acetylsulphadimethoxine-N_1-glucuronide
HPLC of sulphadimethoxine and its N_1-glucuronide, N_4-acetyl and N_4-acetyl-N_1-glucuronide metabolites in human plasma and urine 119
Aminoglycosides
HPLC determination of isepamicin in plasma, urine and dialysate 487
Amyloid A
Purification of human serum amyloid A by anion-exchange fast protein LC 203
Androgen-binding protein
Dihydrotestosterone binding capacity of androgen-binding protein in tissue extract using high-performance size exclusion chromatography 186
Antipyrine
Rapid and simple HPLC determination of saliva antipyrine for routine antipyrine test 296
Apolipoproteins
Purification of human serum amyloid A by anion-exchange fast protein LC 203
Atracurium
Simplified method for the determination of atracurium and laudanosine in pig plasma by HPLC and fluorimetric detection 223
Atrazine
HPLC determination of atrazine in human plasma 569
cis-N-(1-Benzyl-2-methylpyrrolidin-3-yl)-5-chloro-2-methoxy-4-methylaminobenzamide
HPLC determination of *cis*-N-(1-benzyl-2-methylpyrrolidin-3-yl)-5-chloro-2-methoxy-4-methylaminobenzamide in rat blood and brain using electrochemical detection 282
Buspirone
Determination of plasma buspirone by HPLC with coulometric detection 590
Carbonic anhydrase inhibitor
Development of direct stereoselective and non-stereoselective assays in biological fluids for the enantiomers of a thieno[2,3-*b*]thiopyran-2-sulphonamide, a topically effective carbonic anhydrase inhibitor 461
β-Carotene
Endogenous levels of β-carotene in human buccal mucosa cells by reversed-phase HPLC 47
Catecholamines
Simultaneous HPLC determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaestetized normotensive rats 1
Catecholamines
Proposal for the standardization of the calibration method for the assay of plasma catecholamines 194
Choline
Simultaneous HPLC determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaestetized normotensive rats 1
Cocaine
Determination of cocaine and norcocaine in plasma and cell cultures using HPLC 447
Collagen
Determination of collagen and protein turnover by HPLC 383
Cortisol
Semi-automated HPLC method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test 180
Creatine kinase MM
Clinical application of subforms of crea-

tine kinase MM and macro creatine kinases 423

CV-3988
Determination of a platelet activating factor antagonist (CV-3988): methodology and clinical application 515

Cystathionine
Determination of N-monoacetylcystathionine in biological samples using isotachophoresis 550

Cysteine protease inhibitor
HPLC determination of a cysteine protease inhibitor and its ethyl ester in mouse serum and muscle by pre-column fluorescence derivatization 77

Dantrolene
Determination of a dantrolene metabolite, 5-(*p*-nitrophenyl)-2-furoic acid, in plasma and urine by HPLC 575

N-Deacetyl-14-O-demethyllappaconitine
Separation and characterization of the metabolic products of lappaconitine in rat urine by HPLC 109

N-Deacetyl-16-O-demethyllappaconitine
Separation and characterization of the metabolic products of lappaconitine in rat urine by HPLC 109

N-Deacetyl-14,16-O-dimethyllappaconitine
Separation and characterization of the metabolic products of lappaconitine in rat urine by HPLC 109

N-Deacetyllappaconitine
Separation and characterization of the metabolic products of lappaconitine in rat urine by HPLC 109

11-Dehydrothromboxane B_2
11-Dehydrothromboxane B_2: convenient preparation of its stable isotope-labelled analogue for use in quantitative GC–MS 169

N-Demethylencainide
Encainide and metabolites analysis in serum or plasma using a reversed-phase HPLC technique 260

Demethyltauromustine
Determination of tauromustine and its demethylated metabolites in plasma and urine 151

N-Deoxycortisol
Semi-automated HPLC method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test 180

Didemethyltauromustine
Determination of tauromustine and its demethylated metabolites in plasma and urine 151

2′,3′-Dideoxyinosine
HPLC method for analysis of 2′,3′-dideoxyinosine in human body fluids 69

Digoxin, 3,5-dinitrobenzoyl derivative
Electrochemical detection of the 3,5-dinitrobenzoyl derivative of digoxin by HPLC 439

Dihydrotestosterone
Dihydrotestosterone binding capacity of androgen-binding protein in tissue extract using high-performance size exclusion chromatography 186

2,6-Diisopropylphenol
Electrochemical determination of 2,6-diisopropylphenol after HPLC of extracts from serum 293

3,5-Dinitrobenzoyl derivative of digoxin
Electrochemical detection of the 3,5-dinitrobenzoyl derivative of digoxin by HPLC 439

Diphenhydramine
HPLC method for the determination of diphenhydramine in human plasma 597

Ditekiren
Determination of ditekiren, a renin inhibitor peptide, in monkey serum using HPLC with solid-phase extraction 273

Domoid acid
HPLC assay for domoid acid in serum of different species 546

Encainide
Encainide and metabolites analysis in serum or plasma using a reversed-phase HPLC technique 260

Enzymes
Simple isocratic HPLC method for measuring pyridoxine kinase activity in crude biological extracts 25

Enzymes
Clinical application of subforms of crea-

tine kinase MM and macro creatine kinases 423
Erythromycin
Determination of erythromycin base and 2′-acetylerythromycin in human plasma using HPLC with electrochemical detection 475
Etheno adducts
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Ethenoadenine
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Etheno-ADP
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Etheno-AMP
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Etheno-ATP
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Fatty acids
Rapid analysis of non-esterified fatty acids as methyl esters from different biological specimens by GC after one-step esterification 319
Fatty acids
Separation and quantitation of long-chain free fatty acids in human serum by HPLC 331
Fibroblast growth factors
Affinity chromatography of fibroblast growth factors on substituted polystyrene 35
(±)-9-Fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidyl)-5-methyl-1-oxo-1*H*,5*H*-benzo[*i*,*j*]quinolizine-2-carboxylic acid
HPLC procedure for the determination of rat plasma concentrations of a new antibacterial agent, (±)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidyl)-5-methyl-1-oxo-1*H*,5*H*-benzo[*i*,*j*]quinolizine-2-carboxylic acid, for topical use 235
Free fatty acids
Rapid analysis of non-esterified fatty acids as methyl esters from different biological specimen by GC after one-step esterification 319
Free fatty acids
Separation and quantitation of long-chain free fatty acids in human serum by HPLC 331
Gelsolin
Affinity chromatography of human plasma gelsolin with polyphosphate compounds on immobilized Cibracon Blue F3GA 397
Gentamicin
HPLC determination of isepamicin in plasma, urine and dialysate 487
Glimepiride
Simultaneous determination of the sulphonylurea glimepiride and its metabolites in human serum and urine by HPLC after pre-column derivatization 497
Glycerol enol ethers
Simple method for the analysis of glycerol enol ethers derived from plasmalogens in complex lipid mixtures and subsequent determination of the aldehydic components by GC–MS 303
Glycoproteins
Dihydrotestosterone binding capacity of androgen-binding protein in tissue extract using high-performance size exclusion chromatography 186
Glycoproteins
Purification of myelin-associated glycoprotein from calf brain using HPLC 535
Guanidino compounds
Determination of guanidino compounds by anion-exchange chromatography and amperometric detection 355
Hyaluronic acid
Analysis of hyaluronic acid in synovial fluid by reversed-phase LC 530
Hydroxyeicosatetraenoates
Resolution of enantiomers of 19-hydroxyeicosatetraenoate and 18-hydroxyeicosatetraenoate by chiral phase HPLC of naphthoyl ester derivatives 525
5-Hydroxyindoleacetic acid
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367

p-Hydroxyphenyl-phenylhydantoin glucuronide
Direct HPLC analysis of *p*-hydroxyphenyl-phenylhydantoin glucuronide, the final metabolite of phenytoin, in human serum and urine 581
Ibuprofen
Measurement of ibuprofen in human whole blood by reversed-phase ion-paired HPLC using a pH-stable polymeric column 246
Indoleacetic acid
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367
Indolelactic acid
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367
Indolepropionic acid
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367
Indomethacin
Improved procedure for determination of indomethacin in plasma by capillary GC after solid-phase extraction 210
Isepamicin
HPLC determination of isepamicin in plasma, urine and dialysate 487
Kanamycin
HPLC determination of isepamicin in plasma, urine and dialysate 487
Kynurenine
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367
Lappaconitine
Separation and characterization of the metabolic products of lappaconitine in rat urine by HPLC 109
Laudanosine
Simplified method for the determination of atracurium and laudanosine in pig plasma by HPLC and fluorimetric detection 223
Leukotriene E_4
BIO-Fully Automated Sample Treatment HPLC and radioimmunoassay for leukotriene E_4 in human urine from asthmatics 341
Lidocaine
Binding of lidocaine to plasma proteins resolved by HPLC 562
Lipids
Simple method for the analysis of glycerol enol ethers derived from plasmalogens in complex lipid mixtures and subsequent determination of the aldehydic component by GC–MS 303
Macro creatine kinase
Clinical application of subforms of creatine kinase MM and macro creatine kinases 423
Malonaldehyde
Determination of free malonaldehyde by GC with an electron-capture detector 174
Malotilate
Measurement of the formation of a thioglucuronide, a metabolite of malotilate, in rat hepatic microsomes by HPLC 254
Mepindolol
Determination of mepindolol in plasma after transdermal drug delivery by HPLC with electrochemical detection 240
Mercapturic acids
HPLC determination of urinary mercapturic acids in rat 289
3-Methoxy-O-demethylencainide
Encainide and metabolites analysis in serum or plasma using a reversed-phase HPLC technique 260
Metyrapone
Semi-automated HPLC method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test 180
N- Monoacetylcystathionine
Determination of N-monoacetylcystathionine in biological samples using isotachophoresis 550
Myelin-associated glycoprotein
Purification of myelin-associated glycoprotein from calf brain using HPLC 535
Nefopam
Determination of plasma nefopam by LC and electrochemical detection 159
Netilmicin
HPLC determination of isepamicin in plasma, urine and dialysate 487

Nimustine hydrochloride
Rapid and simple method for the determination of nimustine hydrochloride in human blood and brain by HPLC 507
5-(*p*-Nitrophenyl)-2-furoic acid
Determination of a dantrolene metabolite, 5-(*p*-nitrophenyl)-2-furoic acid, in plasma and urine by HPLC 575
Norcocaine
Determination of cocaine and norcocaine in plasma and cell cultures using HPLC 447
Norepinephrine
Simultaneous HPLC determination of norepinephrine, serotonin acetylcholine and their metabolites in the cerebrospinal fluid of anaesthetized normotensive rats 1
Nucleotides
Methods for the isolation and detection of etheno adducts in nucleotide pools in vivo following exposure to ethyl carbamate 407
Phenobarbitone
Metabolic studies with phenobarbitone, primidone and their N-alkyl derivatives: quantification of substrates and metabolites using chemical ionization GC–MS 59
Phenytoin
Direct HPLC analysis of *p*-hydroxyphenyl-phenylhydantoin glucuronide, the final metabolite of phenytoin, in human serum and urine 581
Phospholipids
Solid-phase extraction on silica cartridges as an aid to platelet-activating factor enrichment and analysis 11
Physostigmine
Simultaneous determination of physostigmine and tetrahydroaminoacridine in a transdermal permeation study by HPLC 87
Physostigmine
Sensitive analysis of plasma physostigmine levels using dual-cell electrochemistry in the redox mode 97
Piracetam
Determination of piracetam in serum by GC 556
Plasmalogens
Simple method for the analysis of glycerol enol ethers derived from plasmalogens in complex lipid mixtures and subsequent determination of the aldehydic components by GC–MS 303
Platelet-activating factor
Solid-phase extraction on silica cartridges as an aid to platelet-activating factor enrichment and analysis 11
Platelet activating factor antagonist
Determination of a platelet activating factor antagonist (CV-3988): methodology and clinical application 515
Prilocaine
Determination of prilocaine in human plasma samples using HPLC with dual-electrode electrochemical detection 215
Primidone
Metabolic studies with phenobarbitone, primidone and their N-alkyl derivatives: quantification of substrates and metabolites using chemical ionization GC–MS 59
Propofol
Electrochemical determination of 2,6-diisopropylphenol after HPLC of extracts from serum 293
Protein turnover
Determination of collagen and protein turnover by HPLC 383
Pyridoxine kinase
Simple isocratic HPLC method for measuring pyridoxine kinase activity in crude biological extracts 25
Remoxipride
Determination of remoxipride in plasma and urine by reversed-phase column LC 139
Retinol
Endogenous levels of β-carotene in human buccal mucosa cells by reversed-phase HPLC 47
Retinol palmitate
Endogenous levels of β-carotene in human buccal mucosa cells by reversed-phase HPLC 47
Serotonin
Simultaneous HPLC determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaeasthetized normotensive rats 1
Serotonin
Determination of tryptophan and its metabolites in human plasma and serum by

HPLC with automated sample clean-up system 367

Steroids
Semi-automated HPLC method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test 180

Sulphadimethoxine
HPLC of sulphadimethoxine and its N_1-glucuronide, N_4-acetyl and N_4-acetyl-N_1-glucuronide metabolites in human plasma and urine 119

Sulphadimethoxine-N_1-glucuronide
HPLC of sulphadimethoxine and its N_1-glucuronide, N_4-acetyl and N_4-acetyl-N_1-glucuronide metabolites in human and plasma and urine 119

Sulphide
Determination of sulfide in brain tissue and rumen fluid by ion-interaction reversed-phase HPLC 540

Sulphonylureas
Simultaneous determination of the sulphonylurea glimepiride and its metabolites in human serum and urine by HPLC after pre-column derivatization 497

Taurine
Analysis of taurine in feline plasma and whole blood by LC with fluorimetric detection and confirmation by thermospray MS 375

Tauromustine
Determination of tauromustine and its demethylated metabolites in plasma and urine 151

Tetrahydroaminoacridine
Simultaneous determination of physostigmine and tetrahydroaminoacridine in a transdermal permeation study by HPLC 87

Thieno[2,3-*b*]thiopyran-2-sulphonamide
Development of direct stereoselective and non-stereoselective assays in biological fluids for the enantioners of a thieno[2,3-*b*]thiopyran-2-sulphonamide, a topically effective carbonic anhydrase inhibitor 461

Thioglucuronide
Measurement of the formation of a thioglucuronide, a metabolic of malotilate, in rat hepatic microsomes by HPLC 254

Thromboxane B_2
11-Dehydrothromboxane B_2: convenient preparation of its stable isotope-labelled analogue for use in quantitative GC–MS 169

Tobramycin
HPLC determination of isepamicin in plasma, urine and dialysate 487

Trimetrexate
HPLC assay for trimetrexate in human plasma 266

Tryptophan
Determination of tryptophan and its metabolites in human plasma and serum by HPLC with automated sample clean-up system 367

D-Tubocurarine
Measurement of D-tubocurarine chloride in human urine using solid-phase extraction and reversed-phase HPLC with UV detection 228

Vitamin A
Endogeneous levels of β-carotene in human buccal mucosa cells by reversed-phase HPLC 47